



Older but Not Wiser: the Age-Driven Changes in Neutrophil Responses during Pulmonary Infections

Shaunna R. Simmons,^a Manmeet Bhalla,^a Sydney E. Herring,^a Essi Y. I. Tchalla,^a Elsa N. Bou Ghanem^a

^aDepartment of Microbiology and Immunology, University at Buffalo School of Medicine, Buffalo, New York, USA

ABSTRACT Elderly individuals are at increased risk of life-threatening pulmonary infections. Neutrophils are a key determinant of the disease course of pathogen-induced pneumonia. Optimal host defense balances initial robust pulmonary neutrophil responses to control pathogen numbers, ultimately followed by the resolution of inflammation to prevent pulmonary damage. Recent evidence suggests that phenotypic and functional heterogeneity in neutrophils impacts host resistance to pulmonary pathogens. Apart from their apparent role in innate immunity, neutrophils also orchestrate subsequent adaptive immune responses during infection. Thus, the outcome of pulmonary infections can be shaped by neutrophils. This review summarizes the age-driven impairment of neutrophil responses and the contribution of these cells to the susceptibility of the elderly to pneumonia. We describe how aging is accompanied by changes in neutrophil recruitment, resolution, and function. We discuss how systemic and local changes alter the neutrophil phenotype in aged hosts. We highlight the gap in knowledge of whether these changes in neutrophils also contribute to the decline in adaptive immunity seen with age. We further detail the factors that drive dysregulated neutrophil responses in the elderly and the pathways that may be targeted to rebalance neutrophil activity and boost host resistance to pulmonary infections.

KEYWORDS neutrophils, aging, pneumonia, lung infections, immunosenescence

The population of elderly individuals is on the rise in the United States. Estimates are that 21.7% of the population will be 65 years old or older by 2040 (1). The increased susceptibility to pulmonary infections as we age poses a serious risk to longevity and the quality of life (2). The incidence of community-acquired pneumonia (CAP) is significantly higher in the elderly than in younger adults and is associated with increased mortality, longer time to recovery, and extended hospital stays (3, 4). In fact, approximately 60% of hospital admissions of older adults are estimated to be due to pneumonia (5), and both bacterial and viral infections are the leading causes of CAP in the elderly (6). The Gram-positive bacterium *Streptococcus pneumoniae* (pneumococcus) remains the most frequent bacterial agent of CAP in individuals over the age of 65 years (6), but other bacterial agents associated with CAP in the elderly include *Haemophilus influenzae*, *Moraxella catarrhalis*, *Staphylococcus aureus*, *Mycoplasma pneumoniae*, and *Legionella* species (6). In addition, prior to the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic, the most common viral agents associated with CAP in the elderly were rhinoviruses and influenza viruses, followed by respiratory syncytial virus and parainfluenza viruses (6). Polymicrobial infections are also commonly detected, with bacterial-viral combinations found in >30% of cases (6).

The role of PMNs in pneumonia. Successful control of a pulmonary infection is initially carried out by the host's innate immune response, particularly by polymorphonuclear cells (PMNs), also known as neutrophils (7). PMNs are relatively short-lived cells that are continuously produced in the bone marrow in large numbers (10^{11} /day) (7). Circulating PMNs patrol the body for signs of infection and, in response to local

Citation Simmons SR, Bhalla M, Herring SE, Tchalla EYI, Bou Ghanem EN. 2021. Older but not wiser: the age-driven changes in neutrophil responses during pulmonary infections. *Infect Immun* 89:e00653-20. <https://doi.org/10.1128/IAI.00653-20>.

Editor Anthony R. Richardson, University of Pittsburgh

Copyright © 2021 American Society for Microbiology. All Rights Reserved.

Address correspondence to Elsa N. Bou Ghanem, elsaboug@buffalo.edu.

Accepted manuscript posted online 25 January 2021

Published 17 March 2021




FIG 1 Aging is associated with dysregulated PMN responses. In young hosts, a balanced neutrophil response, where neutrophils migrate to the site of infection, perform antimicrobial functions, and then resolve, is required for proper control and clearance of a pulmonary infection. However, aged hosts experience a decline in PMN function, exacerbated PMN pulmonary influx, and impaired resolution. This unbalanced response results in microbial persistence, tissue damage, and worse disease outcomes.

inflammation, are among the first cells to be recruited to the site of infection to control invading pulmonary pathogens (7) (Fig. 1). Neutropenic individuals, or those with impaired PMN antimicrobial activity, are at increased risk for pneumonia (7–9). In animal models, depletion of PMNs prior to several bacterial (10–13) or viral (14, 15) pulmonary infections results in increased pathogen burdens and heightened mortality. These findings emphasize the importance of PMNs early on in controlling pathogen burdens (Fig. 1). Importantly, however, an exacerbated neutrophilic response can also contribute significantly to lung pathology, because PMNs can inflict major damage on the host (7, 16). In fact, impairment of PMN resolution in the lungs is linked to poorer outcomes, and depletion of these cells after the onset of disease enhances host survival in several models of bacterial and viral pulmonary infections (10, 12, 15). Therefore, successful control of pulmonary infections requires a balance of efficient clearance of invading pathogens by PMNs with the subsequent resolution of these immune cells to prevent tissue damage (Fig. 1).

In this review, we discuss the role of PMNs in the susceptibility of the elderly to pneumonia. We first summarize the age-driven changes in PMN recruitment, resolution, and antimicrobial function (Fig. 1). We highlight the role of PMNs beyond innate immunity and point to the need for further studies to determine whether age-driven changes in PMNs also contribute to the decline in adaptive immunity seen in aging hosts. We conclude by discussing why PMN responses are dysregulated with aging and what pathways may be targeted to rebalance these responses and boost resistance to pulmonary infections (Fig. 2).

AGE-DRIVEN CHANGES IN NEUTROPHIL RESPONSES

Aging is accompanied by persistent low-grade inflammation, known as inflammaging, which, in turn, drives immunosenescence, i.e., the overall decline in immunity seen in the elderly (17). Both phenomena contribute to increased susceptibility of the elderly to pulmonary infections (18, 19). Although PMNs have a shorter life span than other immune cells and are continuously produced in the bone marrow, these innate immune cells also display age-driven impairment (20) (Fig. 1 and 2). However, unlike that of other immune cells, such as naive T cells, the production of PMNs is not diminished in the elderly, who, if healthy, have equivalent or greater numbers of circulating PMNs than




FIG 2 Approaches to rebalancing PMN responses in aged hosts. Several potential pathways and targets have been identified that may boost neutrophil antimicrobial activity, regulate PMN migration, and improve host resistance to infection. Asterisks indicate studies performed only in mice.

younger controls (21–23). Instead, PMN impairment is driven by intrinsic changes, as well as changes in the microenvironment, as discussed below.

In accord with the term “polymorphonuclear,” mature PMNs are characterized by segmented nuclei. These cells differ considerably among species, however, in terms of both their antimicrobial responses and the markers they express. For example, mouse neutrophils express high levels of Ly6G on their surfaces, but Ly6G is not expressed in humans. Rather, human PMNs are mostly characterized by high forward/side scatter and expression of CD16, CD15, or CD66b (24). Since murine models do reflect the age-driven susceptibility of humans to pulmonary infections (15, 25–27), we will focus here on PMN responses from human donors and discuss mouse models of pneumonia that support these findings.

Recruitment. To clear pathogens efficiently, PMNs must reach the site of infection (Fig. 1). In response to microbial challenge, resident pulmonary cells, including alveolar macrophages and lung epithelial cells, produce cytokines and chemokines that recruit PMNs (28). Recruited PMNs move from the circulation across the endothelium, enter the interstitial space, and then cross the lung epithelium into the airways (28). This process involves a complex network of ligand/receptor interactions, as well as cytokine and chemokine receptor signaling (reviewed elsewhere [28]). In aged hosts, general PMN recruitment into the lungs is actually elevated (15, 25, 29–31), but the accuracy of PMN trafficking toward the pulmonary insult is impaired (27, 32, 33) (Fig. 2). With aging, uninfected healthy human donors have elevated PMN levels in the airways; the percentage and number of PMNs in bronchoalveolar lavage fluid (BALF) are significantly higher than those in young controls (34–36). This elevation may be driven by inflamming, since the BALF of elderly donors contains increased levels of cytokines that mediate PMN recruitment, such as interleukin 6 (IL-6) and IL-8 (35, 36). Although very few studies have examined PMN pulmonary influx during pneumonia in elderly individuals, elderly patients with pneumococcal pneumonia exhibit higher percentages of neutrophils in lung biopsy specimens than younger patients (29). While PMNs from healthy elderly volunteers do not display any defects in endothelial adherence (37) or transepithelial migration in response to pneumococcal infection *in vitro* (38), the chemotaxis (directional movement) of PMNs is impaired in aging (32, 33). This impairment is independent of local signaling; instead, it is driven by intrinsic elevated basal levels of phosphoinositide-3-kinase (PI3K) signaling (32). PMNs isolated from healthy elderly donors show reduced chemotaxis toward sputum from *S. pneumoniae* patients relative to that for younger donors (32), and this age-driven impairment in migration is even

more pronounced in circulating PMNs isolated from elderly pneumonia patients, worsening progressively with increased disease severity (33). While chemotaxis toward the stimulus is reduced, chemokinesis, or random movement, of PMNs from elderly donors is not impaired (32). Thus, PMN migration in the elderly is inaccurate, i.e., it does not follow a trajectory toward the stimulus, which may impair the ability of PMNs to efficiently reach the site of infection. This dysregulated movement may contribute to lung damage (Fig. 1), because PMNs from elderly donors release more neutrophil elastase (32, 38). When present in excess, neutrophil elastase can damage pulmonary epithelial cells and vasculature (39), blunt the immune responses of macrophages (40), and impair host resistance to pneumococcal pneumonia, as seen in mouse models (40). Therefore, although aging does not lead to impairment of pulmonary PMN recruitment, PMN colocalization with the pathogen (27), which is necessary for efficient pathogen control and mitigation of bystander damage to tissues, is impaired (Fig. 2).

Resolution. Successful control of a pulmonary infection requires efficient clearance of invading pathogens by immune cells, followed by the resolution of inflammation to prevent tissue damage (7) (Fig. 1). However, aging is associated with an enhanced PMN presence in the lungs in response to pulmonary infections (Fig. 2), observed both in humans (29) and in several mouse models of bacterial and viral pneumonia. PMN numbers in the lungs of old mice are significantly higher than those in young controls during pulmonary infection with *Acinetobacter baumannii* (31), *S. pneumoniae* (25, 30), or influenza A virus (IAV) (15). Following *S. pneumoniae* (10) or IAV infection (41), the numbers of PMNs from the lungs of young mice peak and then resolve over time, but these kinetics are dysregulated with aging, i.e., resolution of pulmonary inflammation is not observed in old mice (15, 25, 30, 41) (Fig. 1). As discussed above, PMN persistence can cause lung damage, whereas experimental depletion of PMNs in the later phase of disease following IAV infection enhances the survival of aged mice without altering the pathogen burden (15). These findings suggest that PMN persistence in the pulmonary space is pathogenic in aged hosts (Fig. 1).

PMN resolution from the lungs is an active process that involves the clearance of apoptotic PMNs present in the pulmonary environment, abrogation of signals that mediate the recruitment of new cells, and production of mediators that negatively regulate PMN influx (42). These factors have been characterized mostly in animal models. One factor that apparently contributes to exacerbated PMN pulmonary presence is the inability of aged hosts to control pathogen burdens. Persistent pathogen detection by resident pulmonary cells leads to the continuous release of inflammatory signals that recruit more cells into the lungs. For example, pneumococcal pneumonia in aged mice results in increased bacterial numbers that correlate with elevated production of inflammatory cytokines that recruit PMNs (25, 30, 43, 44). Similarly, elderly human patients show extended inflammation following pneumococcal infection (45).

PMNs are relatively short-lived cells that undergo spontaneous apoptosis. However, upon PMN entry into inflamed tissue, apoptosis is delayed by inflammatory cytokine signaling in the milieu, allowing PMNs to perform their antibacterial function before death and clearance by resident macrophages. Aging is not accompanied by impaired apoptosis; rather, the ability of inflammatory signals to delay PMN apoptosis is blunted (46, 47). This possibly contributes to the reduced ability of PMNs in aged hosts to control pathogen burdens (Fig. 2) (discussed in the next section). The ability of macrophages to perform efferocytosis, i.e., the engulfment and clearance of apoptotic cells, is impaired with aging (42). In IAV infection of the lungs of aged mice, both the number of alveolar macrophages and their ability to phagocytose apoptotic PMNs are reduced, resulting in impaired PMN resolution (48). One factor, resolvin D1, may contribute to this reduced macrophage efferocytosis. Resolvins belong to a family of specialized proresolving lipid mediators that are crucial for the resolution of inflammation in several organs, including the lungs (49). In mouse models of pulmonary *Escherichia coli* (50) or *Pseudomonas aeruginosa* (51) infection, or in secondary pneumococcal pneumonia (52), resolvin D1 is crucial for host survival and the reduction of PMN influx into the

lungs, as well as for PMN clearance by macrophages. Interestingly, the production of resolvin D1 is reduced in aging (53), and treatment of acute sterile lung injury with this lipid mediator boosts macrophage efferocytosis and ameliorates lung damage in aged mice (54). The role of resolvins in exacerbated PMN responses in elderly individuals during lung infection requires further elucidation.

Extracellular adenosine (EAD) is another regulator of pulmonary PMN influx. Upon tissue damage by infection or a variety of other insults, ATP is thought to leak from damaged cells into the extracellular space, where it can be metabolized to adenosine by the sequential action of two extracellular enzymes: CD39, which converts ATP to AMP, and CD73, which dephosphorylates AMP to EAD (55). EAD can then bind to, and signal via, four G-protein-coupled receptors (56) that are ubiquitously expressed (57). EAD receptor signaling may enhance or diminish acute inflammation, depending on the target receptor, cell type, and/or EAD concentration (56, 57). This pathway has been characterized in several models of sterile inflammation, although its role in pulmonary infections has only recently been appreciated (58–61). In sterile lung injury models, CD73 is important for PMN resolution (62, 63), and EAD production by CD73 has been found both to blunt the transendothelial migration of PMNs into the lung interstitium during pneumococcal pneumonia and to play an essential role in PMN resolution and the prevention of lung damage (10). Similarly, CD73 blunts PMN pulmonary influx during *Mycobacterium tuberculosis* infection (64). Interestingly, CD73 expression on PMNs and EAD production by PMNs in response to *S. pneumoniae* infection are reduced in aged mice (65). This suggests that the EAD pathway is dysregulated during aging and that this dysregulation may contribute to the deficiency in PMN resolution during pulmonary infection.

In summary, several factors have been shown to contribute to the dysregulated PMN responses observed during lung infections of aged hosts (Fig. 2). Elucidating the pathways controlling PMN responses and how they are altered by aging is crucial for devising means to control the regulation of PMN movement and to boost host resistance, as discussed below.

Antimicrobial activity. PMNs kill bacteria by several mechanisms (66). Antimicrobial factors packaged in primary, secondary, and tertiary granules are formed sequentially in PMNs during cellular differentiation in the bone marrow and are released upon encountering a pathogen (67). More-recent findings also show that active transcription is crucial for PMN antimicrobial function, and transcriptional profiling has been used to characterize changes in PMN responses upon influx into the lungs during infection (68, 69). The mechanisms by which PMNs kill microbes include phagocytosis and intracellular killing via the fusion of granules with the phagosome, as well as intracellular production of reactive oxygen species (ROS) (66). PMNs can also kill microbes extracellularly, via degranulation and release of antimicrobial contents, extracellular production of ROS, and the formation of neutrophil extracellular traps (NETs) (66). Many studies have characterized the responses of human PMNs upon *ex vivo* stimulation, revealing that aging is accompanied by overall declines in phagocytosis (70–73), ROS production in response to stimuli (71, 73–75), production of NETs (NETosis) (76), and cytokine production (77, 78) (Fig. 2). However, several studies failed to find age-associated differences in phagocytosis or the oxidative burst (21, 75, 79, 80) (Fig. 2). Fewer studies have examined degranulation; they found either no age-related difference upon stimulation (21, 79) or an increase in the amount of released neutrophil elastase (38), a component of primary granules (Fig. 2). These discrepancies may be driven by several factors, as outlined below.

First, the heterogeneous nature of the elderly, driven by underlying conditions, infections, medications, etc., can confound results when one is studying immune responses (81). To circumvent this issue, the SENIEUR protocol, with very strict inclusion and exclusion criteria, was implemented in the 1980s to assess the immune responses of healthy elderly donors (82). However, only a small percentage (~10%) of elderly people actually satisfy these criteria, and they may not include the individuals at increased risk of infection (81). Interestingly, the immune responses of healthy elderly individuals might contribute to longevity. A study comparing neutrophil

responses across age groups found that the PMN responses of healthy centenarians, i. e., people 100 years of age or older, are comparable to those of young adults with respect to ROS production and phagocytosis, while the same PMN responses are blunted in younger elderly donors (aged 65 to 75 years) (73). In addition, there is a well-known sex-based difference in host susceptibility to infections, as reflected in the incidence of invasive pneumococcal pneumonia, which is higher in males than in females, regardless of age (83, 84). Furthermore, some studies have found that the effect of age on PMN responses is more pronounced in males than in females (73, 85).

A second factor that may confound results is the immune stimulant used. Many studies use either inert beads, bacterial products such as *N*-formylmethionyl-leucyl-phenylalanine (FMLP) formulated peptides and lipopolysaccharides (LPS), or chemicals such as phorbol 12-myristate 13-acetate (PMA) (70–78), while much fewer studies have characterized the age-driven changes in PMN-pathogen interactions (38, 70, 71). While the compounds mentioned above are potent PMN stimulants, they may not always accurately reflect the immune responses to actual pathogens. It is well known that pathogens modulate PMN activities (86), whereas microbial products may not do so accurately. For example, while FMLP inconsistently reveals differences in ROS production across studies (21, 75, 79, 80), ROS production in response to bacterial pathogens, including *S. aureus* (71) and *S. pneumoniae* (87), has consistently been shown to be altered in aging. This response is also pathogen specific. In one study, whereas ROS production by PMNs in response to *S. aureus* was reduced in elderly donors from that in young controls, ROS production in response to *E. coli* was constant across age groups (71). Overall, however, with regard to PMN responses to actual microbes, there is an age-related decline in phagocytosis and intracellular killing of engulfed pathogens, as demonstrated for *Candida albicans* (88), *E. coli*, *S. aureus* (71), and *S. pneumoniae* (70).

The manner in which PMNs “see” a pathogen is a third factor that can influence PMN responses. Serum complement and antibodies can mediate opsonophagocytic uptake and killing of bacteria via complement receptors or antibody (Ab) Fc γ receptors (Fc γ R) expressed on PMNs (89). Several studies have revealed that antibody-mediated PMN responses diminish with age (70, 85). For example, *S. pneumoniae* opsonized with Ab-containing serum from an immunized young donor is killed less efficiently by PMNs from elderly donors than by PMNs from young donors (70). This age-related decline in antibody-mediated killing (85) may be related to a decline in Fc γ RIII (CD16) expression on PMNs (72). In contrast, complement-mediated PMN responses are not impaired with aging. In fact, serum complement activity and the expression of complement receptors on PMNs either remain unchanged or increase with aging (70, 90). Similarly, the ability of PMNs from elderly donors to kill *S. pneumoniae* preopsonized with rabbit complement is actually higher than that for young controls (38). In contrast, opsonization of bacteria with matching serum from each donor consistently demonstrates impaired pathogen killing by PMNs from elderly volunteers (70, 71). These findings suggest that conditions that closely mimic *in vivo* environments (found in the host) more accurately reveal an age-driven decline in pathogen control. In summary, it is important to keep in mind that several aspects of study design can confound the responses of human donors and may mask important age-driven changes in PMN function.

In mouse models, aging is more consistently associated with impaired PMN antibacterial activity. During *S. aureus* skin infections, blunted NETosis is linked to the spread of systemic infection in aged mice (91). In the case of *A. baumannii* pneumonia, aged mice have higher pulmonary CFU counts and lower survival rates than young controls, a finding linked to both reduced release of myeloperoxidase (a component of primary granules) and reduced ROS production by pulmonary PMNs (31). Aging also impairs the ability of murine PMNs to kill engulfed *S. pneumoniae* (65). Importantly, adoptive transfer of PMNs from young controls boosts the resistance of aged mice to pneumococcal pneumonia, highlighting the importance of these cells in host resistance to pulmonary infection (65). While several changes in PMN effector activities in old versus young mice have been described (Fig. 2), the signaling pathways driving these

changes are less well characterized. Elucidating these pathways will be key for improving PMN antimicrobial function, as discussed in “Approaches for reversing dysregulated PMN responses in aging” below.

NEUTROPHIL PHENOTYPES IN AGED HOSTS

Originally thought to be homogeneous, PMNs differ in their expression of surface markers, in their nuclei, and in their density and activity (24, 92, 93). This heterogeneity impacts PMN phenotypes and antibacterial function, as seen in certain disease states and during pulmonary infection (24, 93). The factors that drive PMN heterogeneity remain unclear. During persistent infection and systemic inflammation, the increased demand for PMNs triggers emergency granulopoiesis, characterized by the mobilization of immature precursors from the bone marrow, resulting in the presence of both mature and immature PMNs with distinct characteristics in the circulation (94). Further, direct interaction with bacteria, as well as local reprogramming within the lungs, can alter the phenotypes of PMNs and their ability to kill and clear pathogens (24, 92, 93). However, it remains unclear whether there are bona fide PMN subsets that arise from specific progenitors, and this question is currently being investigated (24, 92, 93). The concept of PMN heterogeneity, the classification of PMN phenotypes, and the question of whether heterogeneity is a result of true subsets, a reflection of different maturation states, or a response to the microenvironment have been extensively reviewed elsewhere (24, 92, 93). Less is currently known about the spectrum of PMN phenotypes during host aging (Fig. 2).

Changes in PMNs driven by infection. Circulating PMNs, which have a higher density than peripheral blood mononuclear cells (PBMCs) (95), are typically isolated via density gradient centrifugation. However, in patients with active *Mycobacterium tuberculosis* infection, lower-density neutrophils (LDNs) that localize with the PBMC fraction are found in the circulation (96). These cells are thought to result from bacterial activation of mature PMNs and to provide an intracellular niche for bacterial replication (96). During pathological conditions, a population of myeloid-derived suppressor cells (MDSCs) that display granulocytic markers and have T cell suppressor functions are also found within the PBMC fraction (97). These granulocytic MDSCs are proposed to be a population of suppressive PMNs that inhibit T cells (97). In humans, granulocytic MDSCs are found in the circulation of *M. tuberculosis* patients with active infection (98) and in cystic fibrosis (CF) patients with *P. aeruginosa* infections (99). In addition, incubation of peripheral blood with bacteria results in the formation of granulocytic MDSCs, suggesting that microbial products can also cause the generation of these cells (99). Importantly, both LDNs and granulocytic MDSCs are associated with more-severe infections and worse disease outcomes (96, 98, 99). In mouse models, granulocytic MDSCs also enhance disease progression during pulmonary infections (100, 101). Aging is associated with an overall expansion of MDSCs (102); however, it is unclear whether granulocytic MDSCs or other LDNs are also altered with aging.

The extracellular-adenosine-producing enzyme CD73 is required for the ability of murine PMNs to kill pneumococci (103). In a mouse model of *S. pneumoniae* lung infection, PMNs recruited into the lungs early on express CD73, which correlates with bacterial control (103). However, as infection progresses, CD73^{low} PMNs accumulate in the lungs and are associated with enhanced bacterial infection (103). CD73^{low} PMNs express lower levels of Ly6G, possibly indicating a less-mature PMN subset. CD73^{low} PMNs are also found in aged mice (65), in which circulating PMNs express lower levels of CD73 than those of young controls, impairing their ability to kill *S. pneumoniae* (65). In humans, PMNs may express CD73 intracellularly (104) rather than on their surfaces (105). Extracellular adenosine signaling is known to play a role in human PMN apoptosis (106), phagocytosis (107), ROS production (108), and degranulation (109), in response to the presence of bacterial products (57). Further, CD73 activity is required for *S. pneumoniae* killing by PMNs in ~60% of young donors (103), suggesting that the expression of this enzyme may also differ among individuals.

Changes in PMNs driven by inflammation. PMNs with distinct phenotypes appear in the circulation during inflammation and can be characterized in whole blood using flow cytometry in the absence of density gradient centrifugation (110). In healthy young volunteers receiving intravenous injections of LPS, three PMN populations, differentiated by their nuclei as well as by CD16 and CD62 ligand (CD62L) expression, appear within hours. The populations consist of segmented mature CD16^{hi} CD62L^{hi} PMNs, immature CD16^{low} CD62L^{hi} band cells, and hypersegmented CD16^{hi} CD62L^{low} cells (110). These cells differ in function; CD16^{hi} CD62L^{low} cells are suppressive PMNs that inhibit T-cell proliferation (110, 111). Further, the populations differ in antibacterial activity: CD16^{low} CD62L^{hi} band cells are efficient at killing *Staphylococcus aureus*, while CD16^{hi} CD62L^{low} cells have impaired antibacterial function. Similar surface changes in circulating PMNs are observed in children with severe viral pneumonia (112), where CD16^{hi} CD62L^{low} PMNs are present in the airways of patients with bacterial coinfections, suggesting that these cells can enter the lungs and possibly play a role in impaired pulmonary control of bacteria. Importantly, CD16^{hi} CD62L^{low} PMN levels are elevated in the circulation of elderly donors relative to those in young controls (74). The presence of these cells is thought to contribute to the impaired bacterial phagocytosis and ROS production detected in total PMNs of elderly donors (74). Inflammation, characterized by increased basal levels of circulating inflammatory cytokines such as IL-6 and tumor necrosis factor alpha (TNF- α) (17, 80), might contribute to the phenotypic and functional changes in PMNs in elderly donors as well. The circulation of frail elderly donors shows an increase in the presence of activated immature band cells, with increased spontaneous ROS production and elevated expression of CD11b. Elevated CD11b expression correlates with increased levels of TNF- α in the sera of donors (87). In mice, genetic abrogation of TNF- α has been shown to prevent this age-driven increase in CD11b expression on PMNs (87). These data suggest that the inflammatory environment in the circulation of elderly donors may play a role in altering the phenotypes and functions of PMNs and thus could be a potential target for combating immunosenescence (Fig. 2).

Changes in PMNs driven by the pulmonary environment. Migration of PMNs from the circulation into tissues results in their activation and potentiation of their antimicrobial activity (7, 113). However, under certain disease conditions, the pulmonary microenvironment can cause aberrant changes in PMNs, resulting in impaired function. This has been described in cystic fibrosis patients (114) and in children with bacterial-viral coinfections (115) and can be modeled *in vitro* using a transwell system seeded with lung epithelial cells. It has been found that migration of PMNs from healthy donors across lung epithelial cells in response to airway fluid supernatant from either cystic fibrosis patients (116) or children with bacterial-viral coinfections (115) resulted in a loss of CD16 (an Fc γ receptor involved in phagocytosis) and an increase in the expression of CD63 (a marker of primary degranulation). These changes are associated with an increase in neutrophil elastase in the airway fluids. Importantly, these migrating PMNs displayed defects in their abilities to kill *P. aeruginosa* (116), *H. influenzae* (115), and *S. aureus* (115). In separate studies (32, 72), PMNs from elderly donors have also been shown to express lower levels of CD16 and elevated levels of CD63. Intriguingly, age-driven changes in the pulmonary microenvironment, including changes in pulmonary epithelial cells and increased levels of neutrophil elastase, are known to occur (35, 36, 117). Thus, age-related changes in the lungs may further contribute to the age-driven PMN impairment seen in pneumonia, suggesting that the effect of the pulmonary microenvironment on PMN antibacterial function in aged hosts is an important area for further investigation.

In summary, changes in the circulation and in the pulmonary environment during aging may contribute to an alteration in PMN responses (Fig. 2). Further characterization of the age-driven changes in PMN phenotype and function, the mechanisms driving these changes, and their link to the susceptibility of the elderly to pneumonia is

needed in order to understand these processes more fully and potentially identify relevant drug targets.

THE ROLE OF NEUTROPHILS BEYOND INNATE IMMUNITY

Apart from their role in innate immunity, recent studies have highlighted the role of PMNs in orchestrating adaptive immune responses (118). PMNs can act as antigen-presenting cells (APCs), directly interact with B cells to induce Ab production, and activate or suppress T-cell responses (118). In some instances, these activities are mediated by PMNs with special characteristics (92). Although aging is accompanied by a well-characterized decline in B- and T-cell responses (119), little is known about how the age-driven alteration of PMN responses may contribute to the decline in adaptive immunity.

Interaction of PMNs with DCs. PMNs can mediate dendritic cell (DC) recruitment and maturation (120). Several bacteria induce murine PMNs to produce chemokines that recruit DCs (120). Human PMNs can also produce chemokines when stimulated *ex vivo*, but interestingly, chemokine production by PMNs from elderly donors is diminished (77, 78). During pulmonary infections, PMNs in the lungs are also important for instructing DCs to migrate to the lung-draining lymph nodes, where they activate T cells (121, 122). For several pulmonary viral infections, including IAV infections, DC activation and trafficking to the draining lymph nodes are diminished in aged mice (41, 123), and early PMN recruitment to the lungs is delayed (41). Therefore, it is possible that changes in PMNs contribute to the impaired DC responses described in aged hosts.

Interaction of PMNs with T cells. PMNs produce a repertoire of chemokines that can recruit T cells to the site of inflammation (124). During IAV infection in mice, PMNs produce chemokine trails that guide CD8⁺ T cells into the trachea (125). Aged mice show a delay in CD8⁺ T-cell recruitment into the lungs (41, 123), but it is unclear whether age-related changes in PMN responses affect the pulmonary recruitment of other immune cells following infection. PMNs can also acquire APC characteristics and present antigen to T cells. In human donors, circulating PMNs acquire DC markers during bacterial infections (126). Upon antigen exposure *ex vivo*, PMNs from young, healthy human donors upregulate major histocompatibility complex class II (MHC-II) and costimulatory molecules (127, 128) and acquire the ability to present IAV antigens to CD4⁺ T cells (127). In mice, PMNs with DC characteristics can acquire and present antigen at the site of infection (129) and have been shown to trigger the proliferation of antigen-specific CD4⁺ T cells during fungal pneumonia (130). Since antigen presentation by professional APCs declines with aging (81, 131), PMNs may thus play an important role in boosting T-cell responses in aged hosts. PMNs can further activate T cells by producing cytokines that can drive T-cell subset differentiation (132), and PMN-derived products, such as NETs, can further prime T cells to respond more efficiently to antigens (133). However, both PMN cytokine production and NETosis have been shown to be blunted in elderly donors (76–78).

On the other hand, PMNs can also suppress T cells, since neutrophil-derived serine proteases, such as neutrophil elastase, can cleave cytokine receptors involved in T-cell activation (134). Interestingly, PMNs from healthy elderly donors have higher neutrophil elastase activity than those from young controls (38); therefore, it is possible that changes in PMNs during aging lead to the suppression of T cells. Further, it has been proposed that granulocytic MDSCs, described above, are a population of suppressive PMNs (97) that produce an arsenal of effectors that inhibit T-cell proliferation (95), including ROS (135) and arginase-1 (136). In some studies, basal ROS production by PMNs has been reported to be elevated in elderly subjects (87), and CD16^{hi} CD62L^{low} cells, thought to be suppressive PMNs (110), are more numerous in the elderly (74). Thus, since aging is associated with a well-characterized decline in T-cell responses (137, 138), it is possible that an impairment of PMN responses that activate T cells or an increase in T-cell-suppressing PMNs contributes to this decline.

Interaction of PMNs with B cells. In human spleen, a subset of PMNs has been found to directly induce Ab production and class switch recombination by B cells in

response to T-cell-independent antigens (139). PMNs play a role in natural Ab responses against pneumococci, and patients with neutropenic disorders have lower levels of Abs to some bacterial strains than healthy controls (139). PMNs that mediate B-cell help are also present in mice (140) and nonhuman primates (141). In mice, PMNs localize with marginal-zone B cells in the spleen and are required for the production of T-cell-independent Abs during systemic *S. pneumoniae* infection (140, 142). Aging is associated with an overall decline in B-cell responses and reduced Ab production and functionality (143, 144). However, the contribution of PMNs to this decline remains unexplored.

Role of PMNs in vaccine responses. In addition to bridging the innate and adaptive responses, PMNs also play an important role in vaccine-mediated protection against pulmonary pathogens (140). PMNs have traditionally been viewed as effectors of vaccine responses, i.e., vaccination triggers Abs, and, as one mechanism of protection, Abs bind to extracellular bacteria and promote their clearance by enhancing uptake and killing by PMNs (70). Evidence that the age-related decline in the intrinsic microbicidal activities of PMNs may contribute to decreased pneumococcal polysaccharide vaccine (PPSV) efficacy in the elderly stems, in part, from the finding that *S. pneumoniae* opsonized with the sera of young donors immunized with PPSV are killed less efficiently by PMNs from elderly donors than by PMNs from young donors. This finding suggests that PMN function is impaired in elderly subjects, even in the presence of specific antibodies (70).

Recent evidence indicates that PMNs also induce vaccine responses in mouse models. In these models, B-helper murine PMNs, described above, are required for Ab production in response to PPSV, which elicits T-cell-independent Ab production (140). Further, PMN depletion at the time of vaccination with T-cell-dependent vaccines in mice also results in the loss of host protection against subsequent challenge (145). Depletion of PMNs during vaccination with a recombinant *M. tuberculosis* protein-based vaccine prevented mice from mounting protective Th1 and Th17 responses and prevented the reduction of pulmonary bacterial burdens in vaccinated mice (146). Similarly, in preclinical studies, PMN depletion in mice at the time of vaccination with Pnevna 13, the pneumococcal polysaccharide conjugate vaccine, resulted in the loss of protective Ab responses and a lack of host protection against subsequent *S. pneumoniae* lung infection (145). Aging leads to defects in both T-cell-dependent and T-cell-independent Ab production following the vaccination of human hosts (143, 144), limiting the efficacy of current vaccines (25, 144). However, the role of PMNs in this phenomenon in human hosts remains an open question. In nonhuman primates, vaccination elicits changes in circulating PMNs (141). Whether PMNs are also activated following human vaccination has not been tested, and it is unclear whether PMNs in the elderly fail to acquire an APC-like provaccine phenotype. Interestingly, although the expression of HLA-DR (human MHC-II) is elevated on PMNs from frail elderly donors relative to that for young controls, the response to vaccines against pulmonary pathogens declines with frailty (144, 147). It is also possible that elderly donors acquire a T-cell-suppressive phenotype that limits immune responses. Therefore, it is important to understand and consider PMN responses in designing vaccines, since PMNs may provide novel targets for boosting responses in vulnerable elderly individuals.

APPROACHES FOR REVERSING DYSREGULATED PMN RESPONSES IN AGING

PMNs are not the only cells with altered responses during aging; immunosenescence affects other innate immune cells, such as macrophages, NK cells, and DCs, as well as cells of the adaptive arm of the immune response, such as B and T cells (19, 20). Therefore, it is unclear whether targeting deficiencies in PMNs alone would be sufficient to boost host resistance to infections. In proof-of-principle experiments, adoptive transfer of PMNs from young, but not aged, mice prior to pneumococcal infection was sufficient to control bacterial numbers and significantly improve disease scores in aged mice (65). In addition, depletion of PMNs a few hours after IAV challenge ameliorated pulmonary damage and boosted the survival of aged mice (15). These findings do

suggest that approaches that target PMN antimicrobial activity, modulate PMN recruitment, or aid in PMN resolution may be successful in boosting the resistance of aged human hosts to pulmonary infection (Fig. 2).

In order to design successful approaches to correct the age-driven defects in PMN response, a better understanding of what drives these changes during host aging is needed. As discussed above, PMN responses in aged hosts are shaped by several factors. The first are intrinsic changes in PMNs, regardless of the environment (32, 65, 70). Second are local changes in the pulmonary microenvironment that influence PMN recruitment (25, 30, 35, 36, 43, 44) and resolution (15, 25, 30, 41) during infections. Third are the systemic changes in inflammatory cytokines that may alter the phenotype of circulating PMNs (17, 80, 87). Both intrinsic and extrinsic factors should be considered in designing approaches to alter PMN responses.

Targeting inflammaging. The low-grade inflammation that accompanies aging is a result of multiple complex factors (17). These include (i) changes in cellular responses that maintain homeostasis, giving rise to the accumulation of misplaced and misfolded self-molecules, including DNA, proteins, and lipids, that are then recognized by pattern recognition receptors, resulting in cytokine production by innate immune cells, (ii) increases in senescent cells, including, but not limited to, T cells, that display a senescence-associated secretory phenotype (SASP), and (iii) age-driven dysbiosis of the microbiota that contributes to, and is driven by, inflammaging (17). Inflammaging is paradoxically associated with the failure of innate immune cells to respond to acute stimuli. In the context of pulmonary infections, this has been well characterized for macrophages: inflammaging impairs the ability of macrophages to respond to *S. pneumoniae*, resulting in an environment permissive to infection (26, 30). Importantly, blocking inflammatory cytokines, such as TNF- α , reverses the age-driven dysregulation of immune responses to *S. pneumoniae* (148, 149). Cytokine production in response to stimuli is diminished in PMNs from elderly donors (77, 78), and as discussed above, TNF- α may affect PMN responses (87). However, the link between inflammaging and PMNs, and the impact of targeting the inflammatory environment on reversing PMN immunosenescence in elderly donors (Fig. 2), need to be elucidated in further detail.

Targeting PMN signaling. Aging is associated with changes in several PMN signaling pathways (150) (Fig. 2). These include an increase in basal levels of activation of mitogen-activated protein kinase (MAPK) pathways (47, 151, 152) and PI3K signaling (32). These changes render PMNs less responsive to acute stimuli and impair their function. Targeting these pathways has been shown to improve outcomes in elderly donors (Fig. 2). In a model of sterile dermal injury, oral administration of a p38 MAPK inhibitor resulted in enhanced PMN clearance in elderly donors (153). Inhibition of PI3K signaling *ex vivo* restores the accurate migration of PMNs from elderly donors to chemotactic signals (32). Treatment of elderly pneumonia patients with high doses of statins, drugs typically used to control cholesterol levels, also restores the migration accuracy of their PMNs in response to *ex vivo* stimuli (33). This effect was observed only in patients with pneumonia that had not progressed to severe disease, suggesting that statins may be beneficial as an early therapeutic intervention. In fact, administration of a high dose of simvastatin to hospitalized pneumonia patients above the age of 55 years has been shown to improve clinical outcomes, since volunteers who received the statin showed increased hospital-free survival (154), suggesting a reduction in pulmonary damage in these patients.

Mechanistic target of rapamycin (mTOR) is a protein kinase signaling complex that plays an important role in immunosenescence (155). In humans, treatment of elderly donors with a combination of mTOR complex 1 inhibitors boosts Ab responses to influenza vaccines and, importantly, decreases the overall incidence of respiratory tract infections (156). Although the specific incidence of pneumonia, a result of infection of the lower respiratory tract, was not reported in these studies, oral administration of rapamycin, an inhibitor of mTOR complex 1, boosted the survival of aged mice upon pulmonary infection with *S. pneumoniae* (157). While mTOR inhibition has been shown to reverse defects in several types of immune cells (158), whether it has an effect on PMN responses remains unclear.

Aging is also accompanied by upstream changes in G-protein-coupled receptor signaling. Extracellular adenosine, produced by CD73 as discussed above, signals via four G-protein receptors: A1, A2A, A2B, and A3. Age-related changes in EAD signaling (159, 160) have been documented in several organs, including the brain (159), pancreas (161), and heart (160). Recent findings suggest that the EAD pathway also plays a role in PMN immunosenescence. Triggering A1 receptor signaling reverses the age-driven impairment in the ability of PMNs to kill *S. pneumoniae* (65) in mice and, importantly, boosts the resistance of aged mice to pulmonary pneumococcal infection (162). In humans, adenosine-based drugs have historically been used to treat heart arrhythmias (163). However, the effect of adenosine-based interventions on PMN immune responses in aging in human volunteers has not been assessed, likely because full A1 receptor agonists have detrimental effects *in vivo* (163). Therefore, it would be useful to devise and test partial agonists, or prodrugs activated by or targeted to PMNs (164).

Nutritional interventions. Vitamin supplementation is a possible approach to preventing respiratory tract infections (Fig. 2). In several studies, supplementation of elderly subjects with high doses (above the recommended intake) of vitamin D₃ (165) or vitamin E (α -tocopherol) (166) has reduced the risk of upper respiratory tract infections. Although these studies did not specifically assess the incidence of pneumonia, both vitamins D (167) and E (38) boost the ability of neutrophils to kill *S. pneumoniae*. In addition, vitamin E reduces PMN migration across pulmonary epithelial cells in response to pneumococci (38). Importantly, oral supplementation of mice with vitamin E reverses the age-associated susceptibility to *S. pneumoniae* lung infection (25). Note that as a potential nutritional intervention, vitamin intake must be well controlled, since excessive intake leads to toxicity (168).

In conclusion, targeting PMN responses to reverse the enhanced susceptibility of the elderly to pulmonary infection is a viable strategy (Fig. 2), and the search for clinical interventions that target PMNs is ongoing. Such therapies will need to balance boosting PMN function with preventing pulmonary damage by these cells.

SUMMARY AND FUTURE PERSPECTIVES

In conclusion, a clear dysregulation of PMN responses in aged hosts contributes to susceptibility to pulmonary infections (Fig. 1). This dysregulation is manifested by inaccurate PMN migration toward the site of infection, a lack of PMN resolution from the lungs, and impaired PMN antimicrobial function (Fig. 2). These changes are likely driven by systemic inflammaging and alterations in the local lung microenvironment, in addition to intrinsic changes in PMNs. Age-driven dysregulated PMN responses impair the innate control of pathogen numbers and may also play a role in subsequent blunted adaptive immune responses. The mechanism(s) by which dysregulation of innate immunity drives the impairment of adaptive memory responses requires additional study; interventions that target systemic inflammation or PMNs directly may be viable approaches for boosting the resistance of the elderly to pulmonary infection (Fig. 2). A better understanding of the systemic changes as well as the intrinsic signaling pathways controlling PMN responses is needed for the design of improved interventions that boost immunity against pulmonary infections in the growing and vulnerable elderly population.

ACKNOWLEDGMENTS

We thank Marcia Osburne for critical feedback on the manuscript.

This work was supported by National Institutes of Health grants R00AG051784 and R21AI145370 to E.N.B.G. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

REFERENCES

1. Wells KB, Schoenbaum M, Unutzer J, Lagomasino IT, Rubenstein LV. 1999. Quality of care for primary care patients with depression in managed care. *Arch Fam Med* 8:529–536. <https://doi.org/10.1001/archfam.8.6.529>.
2. Chong CP, Street PR. 2008. Pneumonia in the elderly: a review of the epidemiology, pathogenesis, microbiology, and clinical features. *South Med J* 101:1141–1145. <https://doi.org/10.1097/SMJ.0b013e318181d5b5>.

3. Stupka JE, Mortensen EM, Anzueto A, Restrepo MI. 2009. Community-acquired pneumonia in elderly patients. *Aging Health* 5:763–774. <https://doi.org/10.2217/ahe.09.74>.
4. Ruiz LA, Espana PP, Gomez A, Bilbao A, Jaca C, Aramburu A, Capelastegui A, Restrepo MI, Zalacain R. 2017. Age-related differences in management and outcomes in hospitalized healthy and well-functioning bacteremic pneumococcal pneumonia patients: a cohort study. *BMC Geriatr* 17:130. <https://doi.org/10.1186/s12877-017-0518-0>.
5. Mouton CP, Bazaldua OV, Pierce B, Espino DV. 2001. Common infections in older adults. *Am Fam Physician* 63:257–268.
6. Wunderink RG, Waterer G. 2017. Advances in the causes and management of community acquired pneumonia in adults. *BMJ* 358:j2471. <https://doi.org/10.1136/bmj.j2471>.
7. Pechous RD. 2017. With friends like these: the complex role of neutrophils in the progression of severe pneumonia. *Front Cell Infect Microbiol* 7:160. <https://doi.org/10.3389/fcimb.2017.00160>.
8. Rolston KV. 2001. The spectrum of pulmonary infections in cancer patients. *Curr Opin Oncol* 13:218–223. <https://doi.org/10.1097/00001622-200107000-00002>.
9. Grudzinska FS, Brodlić M, Scholefield BR, Jackson T, Scott A, Thickett DR, Sapey E. 2020. Neutrophils in community-acquired pneumonia: parallels in dysfunction at the extremes of age. *Thorax* 75:164–171. <https://doi.org/10.1136/thoraxjnl-2018-212826>.
10. Bou Ghanem EN, Clark S, Roggensack SE, McIver SR, Alcaide P, Haydon PG, Leong JM. 2015. Extracellular adenosine protects against *Streptococcus pneumoniae* lung infection by regulating pulmonary neutrophil recruitment. *PLoS Pathog* 11:e1005126. <https://doi.org/10.1371/journal.ppat.1005126>.
11. Garvy BA, Harmsen AG. 1996. The importance of neutrophils in resistance to pneumococcal pneumonia in adult and neonatal mice. *Inflammation* 20:499–512. <https://doi.org/10.1007/BF01487042>.
12. Zhang X, Majlessi L, Deriaud E, Leclerc C, Lo-Man R. 2009. Coactivation of Syk kinase and MyD88 adaptor protein pathways by bacteria promotes regulatory properties of neutrophils. *Immunity* 31:761–771. <https://doi.org/10.1016/j.immuni.2009.09.016>.
13. Tateda K, Moore TA, Deng JC, Newstead MW, Zeng X, Matsukawa A, Swanson MS, Yamaguchi K, Standiford TJ. 2001. Early recruitment of neutrophils determines subsequent T1/T2 host responses in a murine model of *Legionella pneumophila* pneumonia. *J Immunol* 166:3355–3361. <https://doi.org/10.4049/jimmunol.166.5.3355>.
14. Camp JV, Jonsson CB. 2017. A role for neutrophils in viral respiratory disease. *Front Immunol* 8:550. <https://doi.org/10.3389/fimmu.2017.00550>.
15. Kulkarni U, Zemans RL, Smith CA, Wood SC, Deng JC, Goldstein DR. 2019. Excessive neutrophil levels in the lung underlie the age-associated increase in influenza mortality. *Mucosal Immunol* 12:545–554. <https://doi.org/10.1038/s41385-018-0115-3>.
16. Bhowmick R, Maung N, Hurley BP, Bou Ghanem E, Gronert K, McCormick BA, Leong JM. 2013. Systemic disease during *Streptococcus pneumoniae* acute lung infection requires 12-lipoxygenase-dependent inflammation. *J Immunol* 191:5115–5123. <https://doi.org/10.4049/jimmunol.1300522>.
17. Fulop T, Larbi A, Dupuis G, Le Page A, Frost EH, Cohen AA, Witkowski JM, Franceschi C. 2017. Immunosenescence and inflamm-aging as two sides of the same coin: friends or foes? *Front Immunol* 8:1960. <https://doi.org/10.3389/fimmu.2017.01960>.
18. Krone CL, van de Groep K, Trzciński K, Sanders EAM, Bogaert D. 2014. Immunosenescence and pneumococcal disease: an imbalance in host-pathogen interactions. *Lancet Respir Med* 2:141–153. [https://doi.org/10.1016/S2213-2600\(13\)70165-6](https://doi.org/10.1016/S2213-2600(13)70165-6).
19. Kovacs EJ, Boe DM, Boule LA, Curtis BJ. 2017. Inflammaging and the lung. *Clin Geriatr Med* 33:459–471. <https://doi.org/10.1016/j.cger.2017.06.002>.
20. Boe DM, Boule LA, Kovacs EJ. 2017. Innate immune responses in the ageing lung. *Clin Exp Immunol* 187:16–25. <https://doi.org/10.1111/cei.12881>.
21. Corberand JX, Laharrague PF, Fillola G. 1986. Neutrophils of healthy aged humans are normal. *Mech Ageing Dev* 36:57–63. [https://doi.org/10.1016/0047-6374\(86\)90138-7](https://doi.org/10.1016/0047-6374(86)90138-7).
22. Born J, Uthgenannt D, Dodt C, Nunninghoff D, Ringvold E, Wagner T, Fehm HL. 1995. Cytokine production and lymphocyte subpopulations in aged humans. An assessment during nocturnal sleep. *Mech Ageing Dev* 84:113–126. [https://doi.org/10.1016/0047-6374\(95\)01638-4](https://doi.org/10.1016/0047-6374(95)01638-4).
23. Butcher S, Chahel H, Lord JM. 2000. Ageing and the neutrophil: no appetite for killing? *Immunology* 100:411–416. <https://doi.org/10.1046/j.1365-2567.2000.00079.x>.
24. Silvestre-Roig C, Fridlender ZG, Glogauer M, Scapini P. 2019. Neutrophil diversity in health and disease. *Trends Immunol* 40:565–583. <https://doi.org/10.1016/j.it.2019.04.012>.
25. Bou Ghanem EN, Clark S, Du X, Wu D, Camilli A, Leong JM, Meydani SN. 2015. The alpha-tocopherol form of vitamin E reverses age-associated susceptibility to *Streptococcus pneumoniae* lung infection by modulating pulmonary neutrophil recruitment. *J Immunol* 194:1090–1099. <https://doi.org/10.4049/jimmunol.1402401>.
26. Hinojosa E, Boyd AR, Orihuela CJ. 2009. Age-associated inflammation and toll-like receptor dysfunction prime the lungs for pneumococcal pneumonia. *J Infect Dis* 200:546–554. <https://doi.org/10.1086/600870>.
27. Chen MM, Palmer JL, Plackett TP, Deburghraeve CR, Kovacs EJ. 2014. Age-related differences in the neutrophil response to pulmonary *Pseudomonas* infection. *Exp Gerontol* 54:42–46. <https://doi.org/10.1016/j.exger.2013.12.010>.
28. Craig A, Mai J, Cai S, Jeyaseelan S. 2009. Neutrophil recruitment to the lungs during bacterial pneumonia. *Infect Immun* 77:568–575. <https://doi.org/10.1128/IAI.00832-08>.
29. Menter T, Giefing-Kroell C, Grubeck-Loebenstien B, Tzankov A. 2014. Characterization of the inflammatory infiltrate in *Streptococcus pneumoniae* pneumonia in young and elderly patients. *Pathobiology* 81:160–167. <https://doi.org/10.1159/000360165>.
30. Boyd AR, Shivshankar P, Jiang S, Berton MT, Orihuela CJ. 2012. Age-related defects in TLR2 signaling diminish the cytokine response by alveolar macrophages during murine pneumococcal pneumonia. *Exp Gerontol* 47:507–518. <https://doi.org/10.1016/j.exger.2012.04.004>.
31. Gu H, Liu D, Zeng X, Peng LS, Yuan Y, Chen ZF, Zou QM, Shi Y. 2018. Aging exacerbates mortality of *Acinetobacter baumannii* pneumonia and reduces the efficacies of antibiotics and vaccine. *Aging (Albany, NY)* 10:1597–1608. <https://doi.org/10.18632/aging.101495>.
32. Sapey E, Greenwood H, Walton G, Mann E, Love A, Aaronson N, Insall RH, Stockley RA, Lord JM. 2014. Phosphoinositide 3-kinase inhibition restores neutrophil accuracy in the elderly: toward targeted treatments for immunosenescence. *Blood* 123:239–248. <https://doi.org/10.1182/blood-2013-08-519520>.
33. Sapey E, Patel JM, Greenwood HL, Walton GM, Hazeldine J, Sadhra C, Parekh D, Dancer RCA, Nightingale P, Lord JM, Thickett DR. 2017. Pulmonary infections in the elderly lead to impaired neutrophil targeting, which is improved by simvastatin. *Am J Respir Crit Care Med* 196:1325–1336. <https://doi.org/10.1164/rccm.201704-0814OC>.
34. Pignatti P, Ragnoli B, Radaeli A, Moscato G, Malerba M. 2011. Age-related increase of airway neutrophils in older healthy nonsmoking subjects. *Rejuvenation Res* 14:365–370. <https://doi.org/10.1089/rej.2010.1150>.
35. Meyer KC, Ershler W, Rosenthal NS, Lu XG, Peterson K. 1996. Immune dysregulation in the aging human lung. *Am J Respir Crit Care Med* 153:1072–1079. <https://doi.org/10.1164/ajrccm.153.3.8630547>.
36. Meyer KC, Rosenthal NS, Soergel P, Peterson K. 1998. Neutrophils and low-grade inflammation in the seemingly normal aging human lung. *Mech Ageing Dev* 104:169–181. [https://doi.org/10.1016/S0047-6374\(98\)00065-7](https://doi.org/10.1016/S0047-6374(98)00065-7).
37. Nogueira-Neto J, Cardoso AS, Monteiro HP, Fonseca FL, Ramos LR, Junqueira VB, Simão KA. 2016. Basal neutrophil function in human aging: implications in endothelial cell adhesion. *Cell Biol Int* 40:796–802. <https://doi.org/10.1002/cbin.10618>.
38. Bou Ghanem EN, Lee JN, Joma BH, Meydani SN, Leong JM, Panda A. 2017. The alpha-tocopherol form of vitamin E boosts elastase activity of human PMNs and their ability to kill *Streptococcus pneumoniae*. *Front Cell Infect Microbiol* 7:161. <https://doi.org/10.3389/fcimb.2017.00161>.
39. Domon H, Oda M, Maekawa T, Nagai K, Takeda W, Terao Y. 2016. *Streptococcus pneumoniae* disrupts pulmonary immune defence via elastase release following pneumolysin-dependent neutrophil lysis. *Sci Rep* 6:38013. <https://doi.org/10.1038/srep38013>.
40. Domon H, Nagai K, Maekawa T, Oda M, Yonezawa D, Takeda W, Hiyoshi T, Tamura H, Yamaguchi M, Kawabata S, Terao Y. 2018. Neutrophil elastase subverts the immune response by cleaving Toll-like receptors and cytokines in pneumococcal pneumonia. *Front Immunol* 9:732. <https://doi.org/10.3389/fimmu.2018.00732>.
41. Toapanta FR, Ross TM. 2009. Impaired immune responses in the lungs of aged mice following influenza infection. *Respir Res* 10:112. <https://doi.org/10.1186/1465-9921-10-112>.
42. Sendama W. 2020. The effect of ageing on the resolution of inflammation. *Ageing Res Rev* 57:101000. <https://doi.org/10.1016/j.arr.2019.101000>.

43. Williams AE, Jose RJ, Brown JS, Chambers RC. 2015. Enhanced inflammation in aged mice following infection with *Streptococcus pneumoniae* is associated with decreased IL-10 and augmented chemokine production. *Am J Physiol Lung Cell Mol Physiol* 308:L539–L549. <https://doi.org/10.1152/ajplung.00141.2014>.
44. Janesch P, Stulik L, Rouha H, Varga C, Steinhäuser J, Zerbs M, Nagy E. 2018. Age-related changes in the levels and kinetics of pulmonary cytokine and chemokine responses to *Streptococcus pneumoniae* in mouse pneumonia models. *Cytokine* 111:389–397. <https://doi.org/10.1016/j.cyto.2018.09.012>.
45. Bruunsgaard H, Skinhoj P, Qvist J, Pedersen BK. 1999. Elderly humans show prolonged in vivo inflammatory activity during pneumococcal infections. *J Infect Dis* 180:551–554. <https://doi.org/10.1086/314873>.
46. Tortorella C, Simone O, Piazzolla G, Stella I, Cappiello V, Antonaci S. 2006. Role of phosphoinositide 3-kinase and extracellular signal-regulated kinase pathways in granulocyte macrophage-colony-stimulating factor failure to delay Fas-induced neutrophil apoptosis in elderly humans. *J Gerontol A Biol Sci Med Sci* 61:1111–1118. <https://doi.org/10.1093/gerona/61.11.1111>.
47. Fortin CF, Larbi A, Dupuis G, Lesur O, Fulop T, Jr. 2007. GM-CSF activates the Jak/STAT pathway to rescue polymorphonuclear neutrophils from spontaneous apoptosis in young but not elderly individuals. *Biogerontology* 8:173–187. <https://doi.org/10.1007/s10522-006-9067-1>.
48. Wong CK, Smith CA, Sakamoto K, Kaminski N, Koff JL, Goldstein DR. 2017. Aging impairs alveolar macrophage phagocytosis and increases influenza-induced mortality in mice. *J Immunol* 199:1060–1068. <https://doi.org/10.4049/jimmunol.1700397>.
49. Uddin M, Levy BD. 2011. Resolvins: natural agonists for resolution of pulmonary inflammation. *Prog Lipid Res* 50:75–88. <https://doi.org/10.1016/j.plipres.2010.09.002>.
50. Abdulnour RE, Sham HP, Douda DN, Colas RA, Dalli J, Bai Y, Ai X, Serhan CN, Levy BD. 2016. Aspirin-triggered resolvin D1 is produced during self-resolving gram-negative bacterial pneumonia and regulates host immune responses for the resolution of lung inflammation. *Mucosal Immunol* 9:1278–1287. <https://doi.org/10.1038/mi.2015.129>.
51. Codagnone M, Cianci E, Lamolinara A, Mari VC, Nespoli A, Isopi E, Mattosio D, Arita M, Bragonzi A, Iezzi M, Romano M, Recchiuti A. 2018. Resolvin D1 enhances the resolution of lung inflammation caused by long-term *Pseudomonas aeruginosa* infection. *Mucosal Immunol* 11:35–49. <https://doi.org/10.1038/mi.2017.36>.
52. Wang H, Anthony D, Yatmaz S, Wijburg O, Satzke C, Levy B, Vlahos R, Bozinovski S. 2017. Aspirin-triggered resolvin D1 reduces pneumococcal lung infection and inflammation in a viral and bacterial coinfection pneumonia model. *Clin Sci (Lond)* 131:2347–2362. <https://doi.org/10.1042/CS20171006>.
53. Arnardottir HH, Dalli J, Colas RA, Shinohara M, Serhan CN. 2014. Aging delays resolution of acute inflammation in mice: reprogramming the host response with novel nano-proresolving medicines. *J Immunol* 193:4235–4244. <https://doi.org/10.4049/jimmunol.1401313>.
54. Rymut N, Heinz J, Sadhu S, Hosseini Z, Riley CO, Marinello M, Maloney J, MacNamara KC, Spite M, Fredman G. 2020. Resolvin D1 promotes efferocytosis in aging by limiting senescent cell-induced MerTK cleavage. *FASEB J* 34:597–609. <https://doi.org/10.1096/fj.201902126R>.
55. Thompson LF, Eltzschig HK, Ibla JC, Van De Wiele CJ, Resta R, Morote-Garcia JC, Colgan SP. 2004. Crucial role for ecto-5'-nucleotidase (CD73) in vascular leakage during hypoxia. *J Exp Med* 200:1395–1405. <https://doi.org/10.1084/jem.20040915>.
56. Hasko G, Linden J, Cronstein B, Pacher P. 2008. Adenosine receptors: therapeutic aspects for inflammatory and immune diseases. *Nat Rev Drug Discov* 7:759–770. <https://doi.org/10.1038/nrd2638>.
57. Barletta KE, Ley K, Mehrad B. 2012. Regulation of neutrophil function by adenosine. *Arterioscler Thromb Vasc Biol* 32:856–864. <https://doi.org/10.1161/ATVBAHA.111.226845>.
58. Lee JS, Yilmaz O. 2018. Unfolding role of a danger molecule adenosine signaling in modulation of microbial infection and host cell response. *Int J Mol Sci* 19:199. <https://doi.org/10.3390/ijms19010199>.
59. Aeffner F, Woods PS, Davis IC. 2014. Activation of A1-adenosine receptors promotes leukocyte recruitment to the lung and attenuates acute lung injury in mice infected with influenza A/WSN/33 (H1N1) virus. *J Virol* 88:10214–10227. <https://doi.org/10.1128/JVI.01068-14>.
60. Barletta KE, Cagnina RE, Burdick MD, Linden J, Mehrad B. 2012. Adenosine A(2B) receptor deficiency promotes host defenses against Gram-negative bacterial pneumonia. *Am J Respir Crit Care Med* 186:1044–1050. <https://doi.org/10.1164/rccm.201204-0622OC>.
61. Theatre E, Frederix K, Guilmain W, Delierneux C, Lecut C, Bettendorff L, Bours V, Oury C. 2012. Overexpression of CD39 in mouse airways promotes bacteria-induced inflammation. *J Immunol* 189:1966–1974. <https://doi.org/10.4049/jimmunol.1102600>.
62. Eltzschig HK, Thompson LF, Karhausen J, Cotta RJ, Ibla JC, Robson SC, Colgan SP. 2004. Endogenous adenosine produced during hypoxia attenuates neutrophil accumulation: coordination by extracellular nucleotide metabolism. *Blood* 104:3986–3992. <https://doi.org/10.1182/blood-2004-06-2066>.
63. Eckle T, Fullbier L, Wehrmann M, Khoury J, Mittelbronn M, Ibla J, Rosenberger P, Eltzschig HK. 2007. Identification of ectonucleotidases CD39 and CD73 in innate protection during acute lung injury. *J Immunol* 178:8127–8137. <https://doi.org/10.4049/jimmunol.178.12.8127>.
64. Petit-Jentreau L, Jouvion G, Charles P, Majlessi L, Gicquel B, Tailleux L. 2015. Ecto-5'-nucleotidase (CD73) deficiency in *Mycobacterium tuberculosis*-infected mice enhances neutrophil recruitment. *Infect Immun* 83:3666–3674. <https://doi.org/10.1128/IAI.00418-15>.
65. Bhalla M, Simmons SR, Abamonte A, Herring SE, Roggensack SE, Bou Ghanem EN. 2020. Extracellular adenosine signaling reverses the age-driven decline in the ability of neutrophils to kill *Streptococcus pneumoniae*. *Aging Cell* 19:e13218. <https://doi.org/10.1111/accel.13218>.
66. Kolaczowska E, Kubas P. 2013. Neutrophil recruitment and function in health and inflammation. *Nat Rev Immunol* 13:159–175. <https://doi.org/10.1038/nri3399>.
67. Gullberg U, Andersson E, Garwicz D, Lindmark A, Olsson I. 1997. Biosynthesis, processing and sorting of neutrophil proteins: insight into neutrophil granule development. *Eur J Haematol* 58:137–153. <https://doi.org/10.1111/j.1600-0609.1997.tb00940.x>.
68. Gomez JC, Dang H, Kanke M, Hagan RS, Mock JR, Kelada SNP, Sethupathy P, Doerschuk CM. 2017. Predicted effects of observed changes in the mRNA and microRNA transcriptome of lung neutrophils during *S. pneumoniae* pneumonia in mice. *Sci Rep* 7:11258. <https://doi.org/10.1038/s41598-017-11638-7>.
69. Ritchie ND, Evans TJ. 2019. Dual RNA-seq in *Streptococcus pneumoniae* infection reveals compartmentalized neutrophil responses in lung and pleural space. *mSystems* 4:e00216–19. <https://doi.org/10.1128/mSystems.00216-19>.
70. Simell B, Vuorela A, Ekstrom N, Palmu A, Reunanen A, Meri S, Kayhty H, Vakevainen M. 2011. Aging reduces the functionality of anti-pneumococcal antibodies and the killing of *Streptococcus pneumoniae* by neutrophil phagocytosis. *Vaccine* 29:1929–1934. <https://doi.org/10.1016/j.vaccine.2010.12.121>.
71. Wenisch C, Patruta S, Daxböck F, Krause R, Horl W. 2000. Effect of age on human neutrophil function. *J Leukoc Biol* 67:40–45. <https://doi.org/10.1002/jlb.67.1.40>.
72. Butcher SK, Chahal H, Nayak L, Sinclair A, Henriquez NV, Sapey E, O'Mahony D, Lord JM. 2001. Senescence in innate immune responses: reduced neutrophil phagocytic capacity and CD16 expression in elderly humans. *J Leukoc Biol* 70:881–886.
73. Alonso-Fernandez P, Puerto M, Mate I, Ribera JM, de la Fuente M. 2008. Neutrophils of centenarians show function levels similar to those of young adults. *J Am Geriatr Soc* 56:2244–2251. <https://doi.org/10.1111/j.1532-5415.2008.02018.x>.
74. Sauce D, Dong Y, Campillo-Gimenez L, Casulli S, Bayard C, Autran B, Boddart J, Appay V, Elbim C. 2017. Reduced oxidative burst by primed neutrophils in the elderly individuals is associated with increased levels of the CD16^{bright}/CD62L^{dim} immunosuppressive subset. *J Gerontol A Biol Sci Med Sci* 72:163–172. <https://doi.org/10.1093/gerona/glw062>.
75. Biasi D, Carletto A, Dell'Agnola C, Caramaschi P, Montesanti F, Zavateri G, Zeminian S, Bellavite P, Bambara LM. 1996. Neutrophil migration, oxidative metabolism, and adhesion in elderly and young subjects. *Inflammation* 20:673–681. <https://doi.org/10.1007/BF01488803>.
76. Hazeldine J, Harris P, Chapple IL, Grant M, Greenwood H, Livesey A, Sapey E, Lord JM. 2014. Impaired neutrophil extracellular trap formation: a novel defect in the innate immune system of aged individuals. *Aging Cell* 13:690–698. <https://doi.org/10.1111/accel.12222>.
77. Qian F, Guo X, Wang X, Yuan X, Chen S, Malawista SE, Bockenstedt LK, Allore HG, Montgomery RR. 2014. Reduced bioenergetics and toll-like receptor 1 function in human polymorphonuclear leukocytes in aging. *Aging (Albany NY)* 6:131–139. <https://doi.org/10.18632/aging.100642>.
78. Dalboni TM, Abe AE, de Oliveira CE, Lara VS, Campanelli AP, Gasparoto CT, Gasparoto TH. 2013. Activation profile of CXCL8-stimulated neutrophils and aging. *Cytokine* 61:716–719. <https://doi.org/10.1016/j.cyto.2013.01.016>.

79. MacGregor RR, Shalit M. 1990. Neutrophil function in healthy elderly subjects. *J Gerontol* 45:M55–M60. <https://doi.org/10.1093/geronj/45.2.m55>.
80. Ogawa K, Suzuki K, Okutsu M, Yamazaki K, Shinkai S. 2008. The association of elevated reactive oxygen species levels from neutrophils with low-grade inflammation in the elderly. *Immun Ageing* 5:13. <https://doi.org/10.1186/1742-4933-5-13>.
81. Chen WH, Kozlovsky BF, Effros RB, Grubeck-Loebenstein B, Edelman R, Szein MB. 2009. Vaccination in the elderly: an immunological perspective. *Trends Immunol* 30:351–359. <https://doi.org/10.1016/j.it.2009.05.002>.
82. Ligthart GJ, Corberand JX, Fournier C, Galanaud P, Hijmans W, Kennes B, Muller-Hermelink HK, Steinmann GG. 1984. Admission criteria for immunogerontological studies in man: the SENIEUR protocol. *Mech Ageing Dev* 28:47–55. [https://doi.org/10.1016/0047-6374\(84\)90152-0](https://doi.org/10.1016/0047-6374(84)90152-0).
83. Wagenvoort GH, Sanders EA, Vlamincx BJ, de Melker HE, van der Ende A, Knol MJ. 2017. Sex differences in invasive pneumococcal disease and the impact of pneumococcal conjugate vaccination in the Netherlands, 2004 to 2015. *Euro Surveill* 22:30481. <https://doi.org/10.2807/1560-7917.ES.2017.22.10.30481>.
84. Gutierrez F, Masia M, Mirete C, Soldan B, Rodriguez JC, Padilla S, Hernandez I, Royo G, Martin-Hidalgo A. 2006. The influence of age and gender on the population-based incidence of community-acquired pneumonia caused by different microbial pathogens. *J Infect* 53:166–174. <https://doi.org/10.1016/j.jinf.2005.11.006>.
85. Fulop T, Jr, Foris G, Worum I, Leovey A. 1985. Age-dependent alterations of Fc gamma receptor-mediated effector functions of human polymorphonuclear leucocytes. *Clin Exp Immunol* 61:425–432.
86. Kobayashi SD, Malachowa N, DeLeo FR. 2018. Neutrophils and bacterial immune evasion. *J Innate Immun* 10:432–441. <https://doi.org/10.1159/000487756>.
87. Verschoor CP, Loukov D, Naidoo A, Puchta A, Johnstone J, Millar J, Lelic A, Novakowski KE, Dorrington MG, Loeb M, Bramson JL, Bowdish DM. 2015. Circulating TNF and mitochondrial DNA are major determinants of neutrophil phenotype in the advanced-age, frail elderly. *Mol Immunol* 65:148–156. <https://doi.org/10.1016/j.molimm.2015.01.015>.
88. Corberand J, Nguyen F, Laharrague P, Fontanilles AM, Gleyzes B, Gyrard E, Senegas C. 1981. Polymorphonuclear functions and aging in humans. *J Am Geriatr Soc* 29:391–397. <https://doi.org/10.1111/j.1532-5415.1981.tb02376.x>.
89. Esposito AL, Clark CA, Poirier WJ. 1990. An assessment of the factors contributing to the killing of type 3 *Streptococcus pneumoniae* by human polymorphonuclear leukocytes in vitro. *APMIS* 98:111–121. <https://doi.org/10.1111/j.1699-0463.1990.tb01009.x>.
90. Bellavia D, Frada G, Di Franco P, Feo S, Franceschi C, Sansoni P, Brai M. 1999. C4, Bf, C3 allele distribution and complement activity in healthy aged people and centenarians. *J Gerontol A Biol Sci Med Sci* 54: B150–B153. <https://doi.org/10.1093/gerona/54.4.b150>.
91. Tseng CW, Kyme PA, Arruda A, Ramanujan VK, Tawackoli W, Liu GY. 2012. Innate immune dysfunction in aged mice facilitate the systemic dissemination of methicillin-resistant *S. aureus*. *PLoS One* 7:e41454. <https://doi.org/10.1371/journal.pone.0041454>.
92. Rosales C. 2018. Neutrophil: a cell with many roles in inflammation or several cell types? *Front Physiol* 9:113. <https://doi.org/10.3389/fphys.2018.00113>.
93. Deniset JF, Kubes P. 2018. Neutrophil heterogeneity: bona fide subsets or polarization states? *J Leukoc Biol* 103:829–838. <https://doi.org/10.1002/JLB.3RI0917-361R>.
94. Manz MG, Boettcher S. 2014. Emergency granulopoiesis. *Nat Rev Immunol* 14:302–314. <https://doi.org/10.1038/nri3660>.
95. Scapini P, Marini O, Tecchio C, Cassatella MA. 2016. Human neutrophils in the saga of cellular heterogeneity: insights and open questions. *Immunol Rev* 273:48–60. <https://doi.org/10.1111/imr.12448>.
96. Deng Y, Ye J, Luo Q, Huang Z, Peng Y, Xiong G, Guo Y, Jiang H, Li J. 2016. Low-density granulocytes are elevated in mycobacterial infection and associated with the severity of tuberculosis. *PLoS One* 11:e0153567. <https://doi.org/10.1371/journal.pone.0153567>.
97. Pillay J, Tak T, Kamp VM, Koenderman L. 2013. Immune suppression by neutrophils and granulocytic myeloid-derived suppressor cells: similarities and differences. *Cell Mol Life Sci* 70:3813–3827. <https://doi.org/10.1007/s00108-013-1286-4>.
98. Du Plessis N, Loebenberg L, Kriel M, von Groote-Bidlingmaier F, Ribechini E, Loxton AG, van Helden PD, Lutz MB, Walzl G. 2013. Increased frequency of myeloid-derived suppressor cells during active tuberculosis and after recent *Mycobacterium tuberculosis* infection suppresses T-cell function. *Am J Respir Crit Care Med* 188:724–732. <https://doi.org/10.1164/rccm.201302-0249OC>.
99. Rieber N, Brand A, Hector A, Graepler-Mainka U, Ost M, Schafer I, Wecker I, Neri D, Wirth A, Mays L, Zundel S, Fuchs J, Handgretinger R, Stern M, Hogardt M, Doring G, Riethmuller J, Kormann M, Hartl D. 2013. Flagellin induces myeloid-derived suppressor cells: implications for *Pseudomonas aeruginosa* infection in cystic fibrosis lung disease. *J Immunol* 190:1276–1284. <https://doi.org/10.4049/jimmunol.1202144>.
100. Knaul JK, Jorg S, Oberbeck-Mueller D, Heinemann E, Scheuermann L, Brinkmann V, Mollenkopf HJ, Yeremeev V, Kaufmann SH, Dorhoi A. 2014. Lung-residing myeloid-derived suppressors display dual functionality in murine pulmonary tuberculosis. *Am J Respir Crit Care Med* 190:1053–1066. <https://doi.org/10.1164/rccm.201405-0828OC>.
101. Periasamy S, Avram D, McCabe A, MacNamara KC, Sellati TJ, Harton JA. 2016. An immature myeloid/myeloid-suppressor cell response associated with necrotizing inflammation mediates lethal pulmonary tularemia. *PLoS Pathog* 12:e1005517. <https://doi.org/10.1371/journal.ppat.1005517>.
102. Bueno V, Sant'Anna OA, Lord JM. 2014. Ageing and myeloid-derived suppressor cells: possible involvement in immunosenescence and age-related disease. *Age (Dordr)* 36:9729. <https://doi.org/10.1007/s11357-014-9729-x>.
103. Siwapornchai N, Lee JN, Tchalla EYI, Bhalla M, Yeoh JH, Roggensack SE, Leong JM, Bou Ghanem EN. 2020. Extracellular adenosine enhances the ability of PMNs to kill *Streptococcus pneumoniae* by inhibiting IL-10 production. *J Leukoc Biol* 108:867–882. <https://doi.org/10.1002/JLB.4MA0120-115RR>.
104. van Waeg G, Van den Berghe G. 1991. Purine catabolism in polymorphonuclear neutrophils. Phorbol myristate acetate-induced accumulation of adenosine owing to inactivation of extracellularly released adenosine deaminase. *J Clin Invest* 87:305–312. <https://doi.org/10.1172/JCI114987>.
105. Eltzschig HK, Macmanus CF, Colgan SP. 2008. Neutrophils as sources of extracellular nucleotides: functional consequences at the vascular interface. *Trends Cardiovasc Med* 18:103–107. <https://doi.org/10.1016/j.tcm.2008.01.006>.
106. Walker BA, Rocchini C, Boone RH, Ip S, Jacobson MA. 1997. Adenosine A2a receptor activation delays apoptosis in human neutrophils. *J Immunol* 158:2926–2931.
107. Salmon JE, Cronstein BN. 1990. Fc gamma receptor-mediated functions in neutrophils are modulated by adenosine receptor occupancy. A1 receptors are stimulatory and A2 receptors are inhibitory. *J Immunol* 145:2235–2240.
108. Cronstein BN, Daguma L, Nichols D, Hutchison AJ, Williams M. 1990. The adenosine/neutrophil paradox resolved: human neutrophils possess both A1 and A2 receptors that promote chemotaxis and inhibit O₂ generation, respectively. *J Clin Invest* 85:1150–1157. <https://doi.org/10.1172/JCI114547>.
109. Bouma MG, Jeunhomme TM, Boyle DL, Dentener MA, Voitenok NN, van den Wildenberg FA, Buurman WA. 1997. Adenosine inhibits neutrophil degranulation in activated human whole blood: involvement of adenosine A2 and A3 receptors. *J Immunol* 158:5400–5408.
110. Pillay J, Kamp VM, van Hoffen E, Visser T, Tak T, Lammers JW, Ulfman LH, Leenen LP, Pickkers P, Koenderman L. 2012. A subset of neutrophils in human systemic inflammation inhibits T cell responses through Mac-1. *J Clin Invest* 122:327–336. <https://doi.org/10.1172/JCI57990>.
111. Liefeld PHC, Pillay J, Vrsekooop N, Heeres M, Tak T, Kox M, Rooijackers SHM, Kuijpers TW, Pickkers P, Leenen LPH, Koenderman L. 2018. Differential antibacterial control by neutrophil subsets. *Blood Adv* 2:1344–1355. <https://doi.org/10.1182/bloodadvances.2017015578>.
112. Cortjens B, Ingelse SA, Calis JC, Vlaar AP, Koenderman L, Bem RA, van Woensel JB. 2017. Neutrophil subset responses in infants with severe viral respiratory infection. *Clin Immunol* 176:100–106. <https://doi.org/10.1016/j.clim.2016.12.012>.
113. Theilgaard-Monch K, Knudsen S, Follin P, Borregaard N. 2004. The transcriptional activation program of human neutrophils in skin lesions supports their important role in wound healing. *J Immunol* 172:7684–7693. <https://doi.org/10.4049/jimmunol.172.12.7684>.
114. Tirouvanziam R, Gernez Y, Conrad CK, Moss RB, Schrijver I, Dunn CE, Davies ZA, Herzenberg LA, Herzenberg LA. 2008. Profound functional and signaling changes in viable inflammatory neutrophils homing to cystic fibrosis airways. *Proc Natl Acad Sci U S A* 105:4335–4339. <https://doi.org/10.1073/pnas.0712386105>.

115. Grunwell JR, Giacalone VD, Stephenson S, Margaroli C, Dobosh BS, Brown MR, Fitzpatrick AM, Tirouvanziam R. 2019. Neutrophil dysfunction in the airways of children with acute respiratory failure due to lower respiratory tract viral and bacterial coinfections. *Sci Rep* 9:2874. <https://doi.org/10.1038/s41598-019-39726-w>.
116. Forrest OA, Ingersoll SA, Preininger MK, Laval J, Limoli DH, Brown MR, Lee FE, Bedi B, Sadikot RT, Goldberg JB, Tangpricha V, Gaggari A, Tirouvanziam R. 2018. Pathological conditioning of human neutrophils recruited to the airway milieu in cystic fibrosis. *J Leukoc Biol* 104:665–675. <https://doi.org/10.1002/JLB.5HI117-454RR>.
117. Shivshankar P, Boyd AR, Le Saux CJ, Yeh IT, Orihuela CJ. 2011. Cellular senescence increases expression of bacterial ligands in the lungs and is positively correlated with increased susceptibility to pneumococcal pneumonia. *Aging Cell* 10:798–806. <https://doi.org/10.1111/j.1474-9726.2011.00720.x>.
118. Rosales C. 2020. Neutrophils at the crossroads of innate and adaptive immunity. *J Leukoc Biol* 108:377–396. <https://doi.org/10.1002/JLB.4MIR0220-574RR>.
119. Weng NP. 2006. Aging of the immune system: how much can the adaptive immune system adapt? *Immunity* 24:495–499. <https://doi.org/10.1016/j.immuni.2006.05.001>.
120. Schuster S, Hurrell B, Tacchini-Cottier F. 2013. Crosstalk between neutrophils and dendritic cells: a context-dependent process. *J Leukoc Biol* 94:671–675. <https://doi.org/10.1189/jlb.1012540>.
121. Park SJ, Burdick MD, Mehrad B. 2012. Neutrophils mediate maturation and efflux of lung dendritic cells in response to *Aspergillus fumigatus* germ tubes. *Infect Immun* 80:1759–1765. <https://doi.org/10.1128/IAI.00097-12>.
122. Blomgran R, Ernst JD. 2011. Lung neutrophils facilitate activation of naive antigen-specific CD4⁺ T cells during *Mycobacterium tuberculosis* infection. *J Immunol* 186:7110–7119. <https://doi.org/10.4049/jimmunol.1100001>.
123. Zhao J, Zhao J, Legge K, Perlman S. 2011. Age-related increases in PGD (2) expression impair respiratory DC migration, resulting in diminished T cell responses upon respiratory virus infection in mice. *J Clin Invest* 121:4921–4930. <https://doi.org/10.1172/JCI59777>.
124. Mantovani A, Cassatella MA, Costantini C, Jaillon S. 2011. Neutrophils in the activation and regulation of innate and adaptive immunity. *Nat Rev Immunol* 11:519–531. <https://doi.org/10.1038/nri3024>.
125. Lim K, Hyun YM, Lambert-Emo K, Capece T, Bae S, Miller R, Topham DJ, Kim M. 2015. Neutrophil trails flu influenza-specific CD8⁺ T cells in the airways. *Science* 349:aaa4352. <https://doi.org/10.1126/science.aaa4352>.
126. Iking-Konert C, Wagner C, Deneff B, Hug F, Schneider M, Andrassy K, Hansch GM. 2002. Up-regulation of the dendritic cell marker CD83 on polymorphonuclear neutrophils (PMN): divergent expression in acute bacterial infections and chronic inflammatory disease. *Clin Exp Immunol* 130:501–508. <https://doi.org/10.1046/j.1365-2249.2002.02008.x>.
127. Vono M, Lin A, Norrby-Teglund A, Koup RA, Liang F, Lore K. 2017. Neutrophils acquire the capacity for antigen presentation to memory CD4⁺ T cells in vitro and ex vivo. *Blood* 129:1991–2001. <https://doi.org/10.1182/blood-2016-10-744441>.
128. Meinderts SM, Baker G, van Wijk S, Beuger BM, Geissler J, Jansen MH, Saris A, Ten Brinke A, Kuijpers TW, van den Berg TK, van Bruggen R. 2019. Neutrophils acquire antigen-presenting cell features after phagocytosis of IgG-opsonized erythrocytes. *Blood Adv* 3:1761–1773. <https://doi.org/10.1182/bloodadvances.2018028753>.
129. Geng S, Matsushima H, Okamoto T, Yao Y, Lu R, Page K, Blumenthal RM, Ward NL, Miyazaki T, Takashima A. 2013. Emergence, origin, and function of neutrophil-dendritic cell hybrids in experimentally induced inflammatory lesions in mice. *Blood* 121:1690–1700. <https://doi.org/10.1182/blood-2012-07-445197>.
130. Fites JS, Gui M, Kernien JF, Negoro P, Dagher Z, Sykes DB, Nett JE, Mansour MK, Klein BS. 2018. An unappreciated role for neutrophil-DC hybrids in immunity to invasive fungal infections. *PLoS Pathog* 14: e1007073. <https://doi.org/10.1371/journal.ppat.1007073>.
131. Pinti M, Appay V, Campisi J, Frasca D, Fulop T, Sauce D, Larbi A, Weinberger B, Cossarizza A. 2016. Aging of the immune system: focus on inflammation and vaccination. *Eur J Immunol* 46:2286–2301. <https://doi.org/10.1002/eji.201546178>.
132. Kalyan S, Kabelitz D. 2014. When neutrophils meet T cells: beginnings of a tumultuous relationship with underappreciated potential. *Eur J Immunol* 44:627–633. <https://doi.org/10.1002/eji.201344195>.
133. Tillack K, Breiden P, Martin R, Sospedra M. 2012. T lymphocyte priming by neutrophil extracellular traps links innate and adaptive immune responses. *J Immunol* 188:3150–3159. <https://doi.org/10.4049/jimmunol.1103414>.
134. Bank U, Reinhold D, Schneemilch C, Kunz D, Synowitz HJ, Ansoorge S. 1999. Selective proteolytic cleavage of IL-2 receptor and IL-6 receptor ligand binding chains by neutrophil-derived serine proteases at foci of inflammation. *J Interferon Cytokine Res* 19:1277–1287. <https://doi.org/10.1089/107999099312957>.
135. Schmielau J, Finn OJ. 2001. Activated granulocytes and granulocyte-derived hydrogen peroxide are the underlying mechanism of suppression of T-cell function in advanced cancer patients. *Cancer Res* 61:4756–4760.
136. Munder M, Schneider H, Luckner C, Giese T, Langhans CD, Fuentes JM, Kropf P, Mueller I, Kolb A, Modolell M, Ho AD. 2006. Suppression of T-cell functions by human granulocyte arginase. *Blood* 108:1627–1634. <https://doi.org/10.1182/blood-2006-11-010389>.
137. Vinuesa CG, Linterman MA, Yu D, MacLennan IC. 2016. Follicular helper T cells. *Annu Rev Immunol* 34:335–368. <https://doi.org/10.1146/annurev-immunol-041015-055605>.
138. Gustafson CE, Weyand CM, Goronzy JJ. 2018. T follicular helper cell development and functionality in immune ageing. *Clin Sci (Lond)* 132:1925–1935. <https://doi.org/10.1042/CS20171157>.
139. Puga I, Cols M, Barra CM, He B, Cassis L, Gentile M, Comerma L, Chorny A, Shan M, Xu W, Magri G, Knowles DM, Tam W, Chiu A, Busset JB, Serrano S, Lorente JA, Bellosillo B, Lloreta J, Juanpere N, Alameda F, Baro T, de Heredia CD, Toran N, Catala A, Torredell M, Fortuny C, Cusi V, Carreras C, Diaz GA, Blander JM, Farber CM, Silvestri G, Cunningham-Rundles C, Calvillo M, Dufour C, Notarangelo LD, Lougaris V, Plebani A, Casanova JL, Ganai SC, Diefenbach A, Arostegui JI, Juan M, Yague J, Mahlaoui N, Donadieu J, Chen K, Cerutti A. 2011. B cell-helper neutrophils stimulate the diversification and production of immunoglobulin in the marginal zone of the spleen. *Nat Immunol* 13:170–180. <https://doi.org/10.1038/ni.2194>.
140. Chorny A, Casas-Recasens S, Sintès J, Shan M, Polentarutti N, Garcia-Escudero R, Walland AC, Yeiser JR, Cassis L, Carrillo J, Puga I, Cunha C, Bastos H, Rodrigues F, Lacerda JF, Morais A, Dieguez-Gonzalez R, Heeger PS, Salvatori G, Carvalho A, Garcia-Sastre A, Blander JM, Mantovani A, Garlanda C, Cerutti A. 2016. The soluble pattern recognition receptor PTX3 links humoral innate and adaptive immune responses by helping marginal zone B cells. *J Exp Med* 213:2167–2185. <https://doi.org/10.1084/jem.20150282>.
141. Musich T, Rahman MA, Mohanram V, Miller-Novak L, Demberg T, Venzon DJ, Felber BK, Franchini G, Pavlakis GN, Robert-Guroff M. 2018. Neutrophil vaccination dynamics and their capacity to mediate B cell help in rhesus macaques. *J Immunol* 201:2287–2302. <https://doi.org/10.4049/jimmunol.1800677>.
142. Deniset JF, Surewaard BG, Lee WY, Kubas P. 2017. Splenic Ly6G^{high} mature and Ly6G^{int} immature neutrophils contribute to eradication of *S. pneumoniae*. *J Exp Med* 214:1333–1350. <https://doi.org/10.1084/jem.20161621>.
143. Buffa S, Bulati M, Pellicano M, Dunn-Walters DK, Wu YC, Candore G, Vitello S, Caruso C, Colonna-Romano G. 2011. B cell immunosenescence: different features of naive and memory B cells in elderly. *Biogerontology* 12:473–483. <https://doi.org/10.1007/s10522-011-9353-4>.
144. Ridda I, Macintyre CR, Lindley R, Gao Z, Sullivan JS, Yuan FF, McIntyre PB. 2009. Immunological responses to pneumococcal vaccine in frail older people. *Vaccine* 27:1628–1636. <https://doi.org/10.1016/j.vaccine.2008.11.098>.
145. Tchalla EYI, Bhalla M, Wohlfert EA, Bou Ghanem EN. 2020. Neutrophils are required during immunization with the pneumococcal conjugate vaccine for protective antibody responses and host defense against infection. *J Infect Dis* 222:1363–1370. <https://doi.org/10.1093/infdis/jiaa242>.
146. Trentini MM, de Oliveira FM, Kipnis A, Junqueira-Kipnis AP. 2016. The role of neutrophils in the induction of specific Th1 and Th17 during vaccination against tuberculosis. *Front Microbiol* 7:898. <https://doi.org/10.3389/fmicb.2016.00898>.
147. Trzonkowski P, Myśliwska J, Pawelec G, Myśliwski A. 2009. From bench to bedside and back: the SENIEUR protocol and the efficacy of influenza vaccination in the elderly. *Biogerontology* 10:83–94. <https://doi.org/10.1007/s10522-008-9155-5>.
148. Puchta A, Naidoo A, Verschoor CP, Loukov D, Thevaranjan N, Mandur TS, Nguyen PS, Jordana M, Loeb M, Xing Z, Kobzik L, Larche MJ, Bowdish DM. 2016. TNF drives monocyte dysfunction with age and results in impaired

- anti-pneumococcal immunity. *PLoS Pathog* 12:e1005368. <https://doi.org/10.1371/journal.ppat.1005368>.
149. Thevaranjan N, Puchta A, Schulz C, Naidoo A, Szamosi JC, Verschoor CP, Loukov D, Schenck LP, Jury J, Foley KP, Schertzer JD, Larche MJ, Davidson DJ, Verdu EF, Surette MG, Bowdish DME. 2018. Age-associated microbial dysbiosis promotes intestinal permeability, systemic inflammation, and macrophage dysfunction. *Cell Host Microbe* 23:570. <https://doi.org/10.1016/j.chom.2018.03.006>.
 150. Fulop T, Larbi A, Douziech N, Fortin C, Guerard KP, Lesur O, Khalil A, Dupuis G. 2004. Signal transduction and functional changes in neutrophils with aging. *Aging Cell* 3:217–226. <https://doi.org/10.1111/j.1474-9728.2004.00110.x>.
 151. Fortin CF, Larbi A, Lesur O, Douziech N, Fulop T, Jr. 2006. Impairment of SHP-1 down-regulation in the lipid rafts of human neutrophils under GM-CSF stimulation contributes to their age-related, altered functions. *J Leukoc Biol* 79:1061–1072. <https://doi.org/10.1189/jlb.0805481>.
 152. Larbi A, Douziech N, Fortin C, Linteau A, Dupuis G, Fulop T, Jr. 2005. The role of the MAPK pathway alterations in GM-CSF modulated human neutrophil apoptosis with aging. *Immun Ageing* 2:6. <https://doi.org/10.1186/1742-4933-2-6>.
 153. De Maeyer RPH, van de Merwe RC, Louie R, Bracken OV, Devine OP, Goldstein DR, Uddin M, Akbar AN, Gilroy DW. 2020. Blocking elevated p38 MAPK restores efferocytosis and inflammatory resolution in the elderly. *Nat Immunol* 21:615–625. <https://doi.org/10.1038/s41590-020-0646-0>.
 154. Sapey E, Patel JM, Greenwood H, Walton GM, Grudzinska F, Parekh D, Mahida RY, Dancer RCA, Lugg ST, Howells PA, Hazeldine J, Newby P, Scott A, Nightingale P, Hill AT, Thickett DR. 2019. Simvastatin improves neutrophil function and clinical outcomes in pneumonia. A pilot randomized controlled clinical trial. *Am J Respir Crit Care Med* 200:1282–1293. <https://doi.org/10.1164/rccm.201812-2328OC>.
 155. Lopez-Otin C, Blasco MA, Partridge L, Serrano M, Kroemer G. 2013. The hallmarks of aging. *Cell* 153:1194–1217. <https://doi.org/10.1016/j.cell.2013.05.039>.
 156. Mannick JB, Morris M, Hockey HP, Roma G, Beibel M, Kulmatycki K, Watkins M, Shavliakadze T, Zhou W, Quinn D, Glass DJ, Klickstein LB. 2018. TORC1 inhibition enhances immune function and reduces infections in the elderly. *Sci Transl Med* 10:eaaq1564. <https://doi.org/10.1126/scitranslmed.aag1564>.
 157. Hinojosa CA, Mgbemena V, Van Roekel S, Austad SN, Miller RA, Bose S, Orihuela CJ. 2012. Enteric-delivered rapamycin enhances resistance of aged mice to pneumococcal pneumonia through reduced cellular senescence. *Exp Gerontol* 47:958–965. <https://doi.org/10.1016/j.exger.2012.08.013>.
 158. Weichhart T, Hengstschlager M, Linke M. 2015. Regulation of innate immune cell function by mTOR. *Nat Rev Immunol* 15:599–614. <https://doi.org/10.1038/nri3901>.
 159. Mackiewicz M, Nikonova EV, Zimmermann JE, Romer MA, Cater J, Galante RJ, Pack AI. 2006. Age-related changes in adenosine metabolic enzymes in sleep/wake regulatory areas of the brain. *Neurobiol Aging* 27:351–360. <https://doi.org/10.1016/j.neurobiolaging.2005.01.015>.
 160. Willems L, Ashton KJ, Headrick JP. 2005. Adenosine-mediated cardioprotection in the aging myocardium. *Cardiovasc Res* 66:245–255. <https://doi.org/10.1016/j.cardiores.2004.11.008>.
 161. Rolband GC, Furth ED, Staddon JM, Rogus EM, Goldberg AP. 1990. Effects of age and adenosine in the modulation of insulin action on rat adipocyte metabolism. *J Gerontol* 45:B174–B178. <https://doi.org/10.1093/geronj/45.5.b174>.
 162. Bhalla M, Hui Yeoh J, Lamneck C, Herring SE, Tchalla EYI, Heinzinger LR, Leong JM, Bou Ghanem EN. 2020. A1 adenosine receptor signaling reduces *Streptococcus pneumoniae* adherence to pulmonary epithelial cells by targeting expression of platelet-activating factor receptor. *Cell Microbiol* 22:e13141. <https://doi.org/10.1111/cmi.13141>.
 163. Jacobson KA, Tosh DK, Jain S, Gao ZG. 2019. Historical and current adenosine receptor agonists in preclinical and clinical development. *Front Cell Neurosci* 13:124. <https://doi.org/10.3389/fncel.2019.00124>.
 164. Peleli M, Fredholm BB, Sobrevia L, Carlstrom M. 2017. Pharmacological targeting of adenosine receptor signaling. *Mol Aspects Med* 55:4–8. <https://doi.org/10.1016/j.mam.2016.12.002>.
 165. Ginde AA, Blatchford P, Breese K, Zarrabi L, Linnebur SA, Wallace JL, Schwartz RS. 2017. High-dose monthly vitamin D for prevention of acute respiratory infection in older long-term care residents: a randomized clinical trial. *J Am Geriatr Soc* 65:496–503. <https://doi.org/10.1111/jgs.14679>.
 166. Meydani SN, Leka LS, Fine BC, Dallal GE, Keusch GT, Singh MF, Hamer DH. 2004. Vitamin E and respiratory tract infections in elderly nursing home residents: a randomized controlled trial. *JAMA* 292:828–836. <https://doi.org/10.1001/jama.292.7.828>.
 167. Subramanian K, Bergman P, Henriques-Normark B. 2017. Vitamin D promotes pneumococcal killing and modulates inflammatory responses in primary human neutrophils. *J Innate Immun* 9:375–386. <https://doi.org/10.1159/000455969>.
 168. Wu D, Meydani SN. 2008. Age-associated changes in immune and inflammatory responses: impact of vitamin E intervention. *J Leukoc Biol* 84:900–914. <https://doi.org/10.1189/jlb.0108023>.

Shaunna R. Simmons received her B.S. degree in biology from St. Thomas Aquinas College, Sparkill, NY, USA, in 2016 and her M.S. degree in microbiology and immunology from the University at Buffalo, Buffalo, NY, USA, in 2020. Ms. Simmons is currently a Ph.D. student in microbiology and immunology at the University at Buffalo, where she studies the age-related decline in neutrophil function following *Streptococcus pneumoniae* infection.



Manmeet Bhalla received his master's degree in microbiology from Panjab University, Chandigarh, India, in 2007 and then worked as a drug safety specialist for 5 years. He started his Ph.D. studies in microbiology at the University of Otago, Dunedin, New Zealand, in the lab of Keith Ireton, where he worked on host cellular pathways involved in the entry and spread of *Listeria monocytogenes*. His work resulted in a number of publications, and he was awarded the doctorate degree in 2018.



The same year, he joined the Bou Ghanem lab at the University at Buffalo, Buffalo, NY, USA, as a postdoctoral associate to explore the role of the extracellular adenosine pathway in shaping the neutrophil response against *S. pneumoniae* infections. Dr. Bhalla is interested in characterizing age-driven changes in neutrophil functions and investigating how these shape host susceptibility to pneumococcal infections.

Sydney E. Herring obtained her B.Sc. from the Rochester Institute of Technology, Rochester, NY, USA, in biotechnology and molecular bioscience. She is currently a third-year Ph.D. student in the Bou Ghanem lab at the University at Buffalo, Buffalo, NY, USA, training in microbiology and immunology. Ms. Herring's project focuses on the extracellular adenosine pathway and its role in neutrophil ROS production as an antimicrobial mechanism in response to *Streptococcus pneumoniae* infection. Her long-term goal is to work in industry, with a strong interest in autoimmune disease and vaccine development.



Essi Y. I. Tchalla is currently a second-year Ph.D. student in microbiology and immunology at the University at Buffalo (UB) in Buffalo, NY, USA. After receiving her bachelor's degree in biomedical sciences at UB in 2018, she took a research technician position in the Bou Ghanem laboratory. During that time, she started working on elucidating the role of neutrophils during vaccination with the pneumococcal conjugate vaccine. This became her main project when she transitioned to a Ph.D. student in 2019. At the completion of her Ph.D. degree, she hopes to apply her acquired skills in a postdoctoral position at a laboratory that focuses on host-pathogen interactions.



Elsa N. Bou Ghanem, Ph.D., is an assistant professor in the Department of Microbiology and Immunology in the University at Buffalo (UB), Buffalo, NY, USA. She obtained her bachelor's and master's degrees in medical lab science and immunology at the American University of Beirut in her country of origin, Lebanon. She then moved to the United States, where she obtained her Ph.D. from the University of Kentucky, working on innate CD8 T-cell responses to *Listeria monocytogenes* with Dr. Sarah D'Orazio. She did her postdoctoral training with Dr. John Leong at Tufts University, where she became interested in innate immune responses to *Streptococcus pneumoniae*. In 2018, she enthusiastically started her lab at UB, exploring immune senescence, inflammation, and infection. Her lab consists of a group of bright scientists focused on elucidating the role of extracellular adenosine pathways in shaping neutrophil responses and age-driven susceptibility to *S. pneumoniae*.

